Tooth Removed — Simple and Surgical

Adolescents: Ages 12 to less than 17 years

Adults: Ages 17 to less than 65 years

Older Adults: 65+

DAYS OF RECOVERY







PAIN SCALE



		MEDICATION	MAXIMUM DAILY DOSE	PAIN RELIEF	SIDE EFFECTS
First line therapy	PURCHASED OVER THE COUNTER	IBUPROFEN 400 mg	2,400 mg	★★★★ Hours of pain relief	
		NAPROXEN SODIUM 440 mg	1,100 mg	★★★★ Hours of pain relief	
		IBUPROFEN 400 mg plus ACETAMINOPHEN 500 mg	Ibuprofen 2,400 mg Acetaminophen 4,000 mg	$\bigstar \bigstar \bigstar \bigstar$ Hours of pain relief	
		NAPROXEN SODIUM 440 mg plus ACETAMINOPHEN 500 mg	Naproxen Sodium 1,100 mg Acetaminophen 4,000 mg	★★★★ Hours of pain relief	
For extended temporary pain management (Optional)	LOCAL ANESTHETIC	LONG-ACTING LOCAL ANESTHETIC	One time only	★★★★ Hours of pain relief	
If NSAID options above are contraindicated	PURCHASED OVER THE COUNTER	ACETAMINOPHEN 1,000 mg	4,000 mg	★★★ Hours of pain relief	
SURGICAL: If NSAID options above are contraindicated OR postoperative pain control is inadequate	PRESCRIPTION	Tablet 1: ACETAMINOPHEN 325 mg plus either OXYCODONE 5 mg or HYDROCODONE 5-7.5 mg and Tablet 2: ACETAMINOPHEN 325 mg	Acetaminophen 4,000 mg lowest effective dose, fewest tablets, and the shortest duration	★★★ Hours of pain relief	



Extreme CAUTION in adolescents. High risk of misuse or substance use disorder. Caregiver consent is advised.



Potential for severe liver damage



constipation



dizziness



drowsiness







needle



This project was financially supported by the Food and Drug Administration (FDA) of the U.S. Department of Health and Human Services (HHS). The contents are those of the author(s) and do not necessarily represent the official views of, nor an endorsement by, FDA/HHS or the U.S. Government.

These guidelines are intended to help inform clinical decision making by prescribers and patients. They are not intended to be used for the purposes of restricting, limiting, delaying, or denying coverage for, or access to, a prescription issued for a legitimate medical purpose by an individual practitioner acting in the usual course of professional practice.





ADA American Dental Association®